



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

Pyrogenic molecular markers: Linking PAH with BPCA analysis

Wiedemeier, Daniel B ; Brodowski, Sonja ; Wiesenberg, Guido L B

Abstract: Molecular characterization of pyrogenic organic matter (PyOM) is of great interest to understand the formation and behavior of these increasingly abundant materials in the environment. Two molecular marker methods have often been used to characterize and trace PyOM: polycyclic aromatic hydrocarbon (PAH) and benzenepolycarboxylic acid (BPCA) analysis. Since both methods target pyrogenic polycyclic compounds, we investigated the linkages between the two approaches using chars that were produced under controlled conditions. Rye and maize straws and their analogues charred at 300, 400 and 500 °C, respectively, were thus analyzed with both methods. Moreover, we also measured BPCAs directly on the lipid extracts, on which PAHs were analyzed, and on the respective extraction residues, too. Both methods revealed important features of the chars, in particular the increasing degree of aromatic condensation with increasing highest heating temperature (HTT). The overlap between the two methods was identified in the lipid fraction, where the proportion of benzenetricarboxylic acids (B3CAs) correlated with PAH abundance. The results confirmed the validity and complementarity of the two molecular marker methods, which will likely continue to play a crucial role in PyOM research due to the recent developments of compound-specific PAH and BPCA stable carbon ($d^{13}C$) and radiocarbon (^{14}C) isotope methods.

DOI: <https://doi.org/10.1016/j.chemosphere.2014.06.046>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-100894>

Journal Article

Accepted Version

Originally published at:

Wiedemeier, Daniel B; Brodowski, Sonja; Wiesenberg, Guido L B (2015). Pyrogenic molecular markers: Linking PAH with BPCA analysis. *Chemosphere*, 119:432-437.

DOI: <https://doi.org/10.1016/j.chemosphere.2014.06.046>

Pyrogenic molecular markers: Linking PAH with BPCA analysis

Daniel B. Wiedemeier^{*,a}, Sonja Brodowski ^{+,b}, Guido L. B. Wiesenberger^a

^a Department of Geography, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

^b INRES Institute of Crop Science and Resource Conservation, University of Bonn, Katzenburgweg 1, 53115 Bonn, Germany

* Corresponding author: phone: (+41) 44 635 52 28; e-mail: daniel.wiedemeier@geo.uzh.ch

⁺ Deceased

11

12 **Abstract**

13 Molecular characterization of pyrogenic organic matter (PyOM) is of great interest to understand
14 the formation and behavior of these increasingly abundant materials in the environment. Two
15 molecular marker methods have often been used to characterize and trace PyOM: polycyclic
16 aromatic hydrocarbon (PAH) and benzenepolycarboxylic acid (BPCA) analysis. Since both
17 methods target pyrogenic polycyclic compounds, we investigated the linkages between the two
18 approaches using chars that were produced under controlled conditions. Rye and maize straws
19 and their analogues charred at 300, 400 and 500 °C, respectively, were thus analyzed with both
20 methods. Moreover, we also measured BPCAs directly on the lipid extracts, on which PAHs
21 were analyzed, and on the respective extraction residues, too. Both methods revealed important
22 features of the chars, in particular the increasing degree of aromatic condensation with
23 increasing highest heating temperature (HTT). The overlap between the two methods was
24 identified in the lipid fraction, where the proportion of benzenetricarboxylic acids (B3CAs)
25 correlated with PAH abundance. The results confirmed the validity and complementarity of the
26 two molecular marker methods, which will likely continue to play a crucial role in PyOM research
27 due to the recent developments of compound-specific PAH and BPCA stable carbon ($\delta^{13}\text{C}$) and
28 radiocarbon (^{14}C) isotope methods.

29

30 **Keywords**

31 Polycyclic aromatic hydrocarbons; benzene polycarboxylic acids; aromatic condensation; black
32 carbon; biochar; pyrolysis temperature

33

34

34

35 **Highlights**

36

37 - PAH and BPCA marker methods for pyrogenic compounds were compared using chars.

38 - Both methods show increasing aromatic condensation of high temperature chars.

39 - Relationships between the two methods were found in the lipid fraction of the chars.

40 - The findings confirm the validity and theoretical assumptions of both methods.

41 - The methods complement each other in pyrogenic carbon research.

42

43

44

45

46 **1. Introduction**

47 Pyrogenic organic matter (PyOM) is ubiquitous in the environment (Schmidt and Noack, 2000;
48 Schmidt et al., 2001; Masiello, 2004) and relatively stable against degradation, making it part of
49 the slow-cycling C pool (Marschner et al., 2008; Kuzyakov et al., 2014). PyOM abundance is
50 thought to increase globally with increasing wildfire occurrence in the 21st century (Flannigan et
51 al., 2013; Kelly et al., 2013) and as a result of industrial combustion processes and traffic (Bond
52 et al., 2013). Moreover, it is increasingly and intentionally produced in anthropogenic biochar-
53 systems, which have been proposed as a strategy for green energy production, C sequestration,
54 and soil improvement (Lehmann and Joseph, 2009; Meyer et al., 2011). With increasing PyOM
55 production and the awareness of its importance for the C cycle, attention has been drawn to
56 characterize these materials accurately across different disciplines (Simoneit, 2002; Hammes et
57 al., 2007; McBeath et al., 2011). Molecular marker methods are one technique providing the
58 opportunity to characterize PyOM. They target specifically the molecules that are produced
59 during the combustion processes. Two different approaches are often used that both focus on
60 the polycyclic structures typical for pyrogenic materials: i) polycyclic aromatic hydrocarbon (PAH)
61 analysis and ii) the benzenepolycarboxylic acid (BPCA) method.

62 PAHs are small polycyclic aromatic compounds that have a long history as tracers for
63 combustion products (Simoneit et al., 1999; Simoneit, 2002; Denis et al., 2012). They proved to
64 be particularly useful for the source apportionment of combustion residues in soils and
65 sediments, for example distinguishing between contributions of biomass burning and fossil fuel
66 (Oros and Simoneit, 2001a, b; Yunker et al., 2002; Bucheli et al., 2004). High concentrations of
67 PAHs are of environmental concern due to their mutagenic and carcinogenic properties (Boffetta
68 et al., 1997; Wilcke, 2000). Thus, PAHs have recently also been heavily investigated in the
69 context of biochar applications (Hale et al., 2012; Hilber et al., 2012; Keiluweit et al., 2012;

Oleszczuk et al., 2013; Quilliam et al., 2013). After extraction with the lipid fraction from bulk PyOM (Colmsjö, 1998; Fabbri et al., 2013), they are routinely quantified with gas or liquid chromatography.

BPCAs, on the other hand, present a workaround to quantify the abundant polycyclic PyOM structures that are larger and more complex than PAHs. These more condensed aromatic C phases cannot be quantified directly as polymers using chromatographic methods. In order to analyze such pyrogenic structures, PyOM is digested with nitric acid under high temperature and pressure, which breaks the polycyclic compounds down into individual BPCAs, amenable to gas or liquid chromatography (Glaser et al., 1998; Brodowski et al., 2005; Wiedemeier et al., 2013). Pyrogenic C is thus assessed on a molecular level and can be used to estimate PyOM abundance in environmental compartments such as soils and sediments. The technique has yielded valuable insights into the slow cycling of PyOM in the pedosphere (Hammes et al., 2008; Rodionov et al., 2010) and has been helpful to illuminate the pathways of PyOM into sedimentary systems, where PyOM accounts for a quantitatively important fraction of the carbon sink (Guggenberger et al., 2008; Sánchez-García et al., 2013). The BPCA method additionally reveals information about the aromaticity, aromatic condensation and charring temperature of the analyzed PyOM when relative yields of benzenetri-, benzenetetra-, benzenepenta- and benzenhexacarboxylic acids (B3-, B4-, B5- and B6CA) are compared (Dittmar, 2008; Schneider et al., 2011; Wiedemeier et al., 2014). These qualitative parameters are indicative for the stability of PyOM in the environment and can also help to produce suitably engineered biochars (Harvey et al., 2012a).

In this study, we investigated the link between the two molecular marker approaches. Despite having different analysis windows, both methods target a part of the characteristic PyOM polycyclic aromatic features that become increasingly condensed with higher charring temperatures (McBeath et al., 2011). The aim was to test if the conceptually feasible overlap

95 between the small aromatic PAH moieties and the larger, condensed polycyclic structures, as
96 indicated by BPCAs, can be assessed with the two methods when applied to real pyrogenic
97 sample materials. We thus measured PAHs and BPCAs on two different straws (rye and maize)
98 and their corresponding chars with highest heating temperatures (HTT) of 300, 400 and 500 °C,
99 respectively. Moreover, we also analyzed BPCAs on the same lipid extracts, on which PAHs
100 were measured, and determined BPCAs on the resulting extraction residues, too.

101

102 **2. Materials and methods**

103 Chars were produced by heating rye (*Secale cereal* L.) and maize (*Zea mays* L.) straw samples
104 for 2 hours in a pre-heated muffle furnace at 300, 400 and 500 °C, respectively (Rennert et al.,
105 2008). Al foil was used to limit oxidation during the charring process. Straw and char samples
106 were subsequently milled to fine powder before chemical analysis.

107 Organic carbon contents were measured using elemental analysis (Leco CS 225,
108 Mönchengladbach Germany). Lipids from all samples were extracted using the accelerated
109 solvent extraction method (Dionex ASE 200, Sunnyvale, CA) with CH₂Cl₂/CH₃OH (93/7; v/v) as
110 described by Wiesenberg et al. (2004; 2008; 2009). Samples were sequentially extracted for 20
111 min in two steps at 5×10^6 Pa and 75 °C and 140 °C, respectively. Both extracts were combined
112 thereafter and cleaned using solid phase extraction columns fitted with KOH coated silica gel
113 (Wiesenberg et al., 2008). Aromatic hydrocarbons were then measured after sequential
114 chromatographic separation (Radke et al., 1980) on GC/MS (HP 5890 Series II GC and HP
115 5989A mass spectrometer, Palo Alto, CA). Detailed data from the PAH analysis of the samples
116 were reported by Wiesenberg et al. (2009), while we here used specific statistics of these results
117 with the focus to assess the linkages between the PAH and BPCA method.

118 BPCAs were measured on bulk materials (the two straws and all their charred analogues) as
119 well as on the respective products from the lipid analysis (the lipid extracts and the lipid
120 extraction residues). The lipid extracts of the 500 °C chars did not yield enough material to be
121 amenable for BPCA analysis. BPCAs were analyzed following the procedure reported by
122 Brodowski et al. (2005), i.e. bulk samples, lipid extracts and lipid extraction residues all
123 underwent the same digestion and analysis steps. Method-inherent underestimation of PyOM
124 was not corrected by any conversion factor (Schneider et al., 2010). For consistency purposes,
125 we used gas chromatography protocols for both PAH and BPCA quantification. However, both

126 methods have recently been adapted to liquid chromatography, which offers an interesting
127 alternative for the assessment of these markers (Schneider et al., 2011), particularly in
128 environmental samples (Denis et al., 2012; Wiedemeier et al., 2013).

129 Data analysis was conducted with the statistical software R (R, 2011). Changes in the
130 distribution of individual BPCAs between the bulk samples, the lipid extracts and the lipid
131 extraction residues were analyzed with the chi-square goodness of fit test. Relationships
132 between BPCAs and PAHs were analyzed with linear regression.

133

134 3. Results and discussion

135 PAHs were undetectable in the straw materials, present in chars with HTT of 300 °C and highly
136 abundant in chars that were pyrolyzed between 400 and 500 °C (Figure 1; top). This is in line
137 with previous studies (Brown et al., 2006; Kloss et al., 2012) and indicates the potentially
138 problematic use of low-temperature biochars as a soil amendment due to their usually high
139 content of harmful PAHs (Keiluweit et al., 2012). Moreover, a distinct pattern between the
140 proportion of PAHs with different ring sizes and the pyrolysis temperature was observed: The
141 proportion of small PAHs (3-ring) decreased while the contribution of higher molecular weight
142 PAHs (4-6 ring) increased with increasing temperature (Figure 1; bottom). Similar trends have
143 been observed in other studies (McGrath et al., 2003; Brown et al., 2006; Keiluweit et al., 2012)
144 and point to the increasingly condensed nature of the solid PyOM residue (the char) while losses
145 of the smaller, more volatile PAHs occur with increasing charring temperature (McGrath et al.,
146 2003).

147 The BPCA method detects these large, condensed polycyclic structures of non-volatile PyOM
148 and consequently showed increasing amounts of BPCA quantity with increasing temperature in
149 the bulk char samples (Figure 2; black bars). Besides the total BPCA yields normalized to bulk
150 weight, also the proportion of BPCA normalized to organic carbon increased with temperature.
151 The latter is a measure for the aromaticity of charred materials (McBeath et al., 2011) and is
152 consistently higher for rye-derived chars than for the maize-based char samples (Figure 2; upper
153 number). This difference in aromaticity was partially reflected by the data obtained from the PAH
154 analysis, as the rye chars showed higher proportions of the larger PAHs than the maize chars
155 (Figure 1). However, the relationship between PAH ring sizes and aromaticity of PyOM are
156 complex and not entirely resolved yet (Wiedemeier et al., 2014). The BPCA analysis furthermore
157 yields a measure for the degree of aromatic condensation with the ratio of B6CAs to total BPCA

content (Schneider et al., 2011). The aromatic condensation was very similar for both, maize and rye chars, at each respective charring temperature and comparable to previously reported values for a variety of charred materials (Schneider et al., 2011; Wiedemeier et al., 2014). It indicates that temperature is the main factor controlling condensation and that an assessment of the aromatic condensation in PyOM is possibly useful to estimate its pyrolysis temperature. Moreover, the degree of aromatic condensation in PyOM has been directly associated to its stability against degradation in the environment (Harvey et al., 2012b; Wurster et al., 2013; Fang et al., 2014). Our results in combination with complementary research (Wiedemeier et al., 2014) therefore support the assumption that the C sequestration potential of charred lignocellulose material is largely determined by its pyrolysis temperature (Bruun et al., 2011; Fang et al., 2014).

When lipid extraction was applied on the bulk straw and char samples, a part of the polycyclic PyOM structures was extracted with it, too, while another part remained in the extraction residue, as shown by the BPCA analysis in Figure 2 (white and striped bars, respectively). The lipid extract always comprised a much smaller part (< 10 % of bulk dry weight) than the extraction residue and became smaller for chars pyrolyzed at higher temperatures, following a trend, which has been reported for other chars, too (Wiesenberg et al., 2009; Wiedemeier et al., 2014). However, the proportion of lipid extracted PyOM increased with charring temperature (Figure 2; white bars). This is explained by the increased availability of these polycyclic structures and the concomitant decrease of other lipids with higher HTT (Wiesenberg and Brodowski, 2007; Wiesenberg et al., 2009). At the same time, the much larger (> 90 % of bulk dry weight) lipid extraction residue comprised a similar proportion of BPCA-detectable PyOM as the lipid extract, also increasing with temperature (Figure 2; striped bars).

Having a closer look at the contributions of the individual BPCAs that make up the above reported BPCA contents can reveal further qualitative information about the PyOM found in bulk samples, lipid extracts and lipid extraction residues (Figure 3). In particular, the size of the PyOM

polycyclic clusters can be inferred, as the more carboxylated B6CA and B5CA must derive from larger polycyclic clusters than the less carboxylated B4- and B3CAs (Hammes et al., 2008; Schneider et al., 2010). Since the extraction residue inherited most of the bulk PyOM by mass (> 90 % of bulk dry weight), it is not surprising that it showed almost the same BPCA pattern like the bulk chars, dominated by B6- and B5CAs (Figure 3). Bulk chars and the lipid extraction residues thus mostly consisted of relatively large, condensed polycyclic sheets, comprising probably the most stable aromatic phases of the chars (Franklin, 1951; Cohen-Ofri et al., 2006; Keiluweit et al., 2010). In contrast, the lipid extractable BPCAs of both rye and maize chars across all pyrolysis temperatures were mostly composed of smaller polycyclic compounds, as indicated by the predominance of B3- and B4CAs. The larger polycyclic clusters that prevailed in the bulk samples (57 – 67 % of B5- and B6CA) thus remained in the extraction residue (57 – 64 % of B5- and B6CA) while small polycyclic compounds were preferentially released during the lipid extraction (51 – 59 % of B3- and B4CA).

The overlap between the PAH and the BPCA method consequently concerns mostly the smallest BPCA-detectable polycyclic clusters. When BPCA analysis was performed on the lipid extracts for direct comparison, we found significant logarithmic relationships ($p < 0.05$, $R^2 > 0.9$) between the proportion of B3CAs and the abundance of most PAHs that were considered in this study (Figure 4). The relationship was independent of the molecular weight of the PAHs and was equally valid for total PAH abundance. However, the proportion of B3CA was the only direct predictor of PAH abundance, while proportions of more carboxylated BPCAs in the lipid fraction showed more complicated relationships with PAH abundance (cf. supplementary material). This is in line with theoretical considerations (Glaser et al., 1998; Dittmar, 2008), after which the majority of the abundant small PAHs (cf. Figure 1) are prone to yield B3- or B4CAs and only less abundant, larger PAH species, such as benzo[a]pyrene or larger condensed polycyclic sheets, can be oxidized into more carboxylated BPCAs. It is unfortunate that the 500 °C chars did not yield enough lipid extract for BPCA analysis, which is why the comparison of the two methods

could not be conducted directly on the lipid extracts for the highest HTT of this study. However, as condensation of bulk chars increased (cf. Figure 2) but extractability of B5- and B6CA decreased with higher temperature (cf. Figure 3), a similar B3CA proportion can be expected in the 500 °C as in the 400 °C lipid extracts, correlating with the still high PAH abundances at 500 °C (cf. Figure 1). Our findings confirm that the analysis windows of the PAH and the BPCA method overlap each other within the lipid fraction. Although the linkage between the two molecular marker methods has been shown for pure chemical standards (Dittmar, 2008), to the best of our knowledge, it is the first time that the theory-based relationship was established empirically on real char samples. However, it is worth noting that no relationship could be established when both methods were applied directly on the bulk chars according to their normal protocol, as the BPCA analysis was then not limited to the lipid fraction only.

This study highlighted the similarities and differences of the two most common molecular marker methods for PyOM. Despite having minimally overlapping analysis windows and traditionally serving different research purposes, they are both highly informative for PyOM characterization as they provide valuable information about its molecular composition that is strongly linked to combustion conditions. We could demonstrate that the two molecular marker approaches find common ground when the lipid fraction is considered, which confirms the validity and theoretical assumptions of both methods. Large benefits can derive from the simultaneous application of the PAH and the BPCA method to relevant environmental sample materials. The combination of both methods can lead to a better source apportionment of PyOM in soils and sediments and illuminate the pathways and fluxes of differently sized and pyrolysed PyOM through environmental compartments. Even more information about PyOM in the environment can be retrieved from compound-specific isotopic analysis of both PAH and BPCA molecular markers (Ziolkowski and Druffel, 2010; Xu et al., 2012; Coppola et al., 2013; Slater et al., 2013; Gierga et al., 2014), which is why it is crucial to understand the here presented relationships between these two approaches.

235

236

236

237 **Acknowledgements**

238 Thilo Rennert (University of Hohenheim, Germany) is thanked for preparing and providing
239 charred samples. We gratefully acknowledge financial support of the current study by the Swiss
240 National Science Foundation under contract 134452 and the German Research Foundation
241 under contract WI 2810 / 9-1.

242

243

Figures





Figure 1.


Absolute (top) and relative (bottom) abundance of PAHs ($n = 1$) changed with highest heating temperature (HTT). Small, 3-ring PAHs were most abundant at HTT of 300 °C and decreased with higher HTT while 4-ring PAHs became more abundant with higher HTT. The larger PAHs (5- and 6-ring) contributed a much smaller proportion to total PAH (cf. y-axis scale is stretched by a factor of 10) and their abundance increased with higher HTT, consistent with an increasing degree of aromatic condensation as revealed by BPCAs (cf. Fig. 2).



257

258 Figure 2.

259 PyOM measured as BPCAs in the bulk materials (maize/rye straw and chars), the respective
 260 lipid extracts and the remaining lipid extraction residues (n = 2 - 5, error bars indicate SE). The
 261 aromaticity and degree of aromatic condensation of the bulk samples increased with increasing
 262 HTT. Likewise, the PyOM content in the lipid extracts and in the lipid extraction residues
 263 increased with increasing HTT. The lipid extracts of the chars produced at 500 °C did not yield
 264 sufficient sample amounts for BPCA analysis.




266

267 Figure 3.

268 BPCA signatures of bulk chars, their lipid extracts and the corresponding lipid extraction
 269 residues ($n = 2 - 5$, $SE \leq 3$ percentage points). Larger PyOM polycyclic structures, indicated by
 270 B5CA and B6CA, were retained in the lipid extraction residues (ns = no significant difference in
 271 BPCA distribution between lipid extraction residue and respective bulk sample analogue). B3CA
 272 and B4CA, indicative of smaller polycyclic compounds, were preferentially extracted and
 273 accounted for more than 50 % of all BPCAs in the lipid extracts (asterisks denote significance of
 274 difference in BPCA distribution between lipid extract and respective bulk sample analogue with *
 275 $p < 0.05$; ** $p < 0.01$ and *** $p < 0.001$).

276



277

278 Figure 4.

279 The proportion of B3CA in the lipid extracts ($n = 2$, $SE \leq 1.4$ percentage points) correlated with
280 PAH abundance ($n = 1$). Data is shown for three different PAH sizes and is indicative for all
281 PAHs considered in this study as well as for total PAH abundance. The correlation confirms that
282 the measurement window of PAH analysis overlaps with B3CA analysis in the lipid extracts while
283 no correlation could be found between PAHs and more carboxylated BPCAs, indicative of larger
284 polycyclic clusters that would escape PAH analysis.

285

286

287 **References**

- 288 Boffetta, P., Jourenkova, N., Gustavsson, P., 1997. Cancer risk from occupational and
289 environmental exposure to polycyclic aromatic hydrocarbons. *Cancer Causes & Control* 8, 444-
290 472.
- 291 Bond, T.C., Doherty, S.J., Fahey, D.W., Forster, P.M., Berntsen, T., Deangelo, B.J., Flanner,
292 M.G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P.K., Sarofim, M.C., Schultz,
293 M.G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S.K.,
294 Hopke, P.K., Jacobson, M.Z., Kaiser, J.W., Klimont, Z., Lohmann, U., Schwarz, J.P., Shindell,
295 D., Storelvmo, T., Warren, S.G., Zender, C.S., 2013. Bounding the role of black carbon in the
296 climate system: A scientific assessment. *Journal of Geophysical Research D: Atmospheres* 118,
297 5380-5552.
- 298 Brodowski, S., Rodionov, A., Haumaier, L., Glaser, B., Amelung, W., 2005. Revised black
299 carbon assessment using benzene polycarboxylic acids. *Organic Geochemistry* 36, 1299-1310.
- 300 Brown, R.A., Kercher, A.K., Nguyen, T.H., Nagle, D.C., Ball, W.P., 2006. Production and
301 characterization of synthetic wood chars for use as surrogates for natural sorbents. *Organic*
302 *Geochemistry* 37, 321-333.
- 303 Bruun, E.W., Hauggaard-Nielsen, H., Ibrahim, N., Egsgaard, H., Ambus, P., Jensen, P.A., Dam-
304 Johansen, K., 2011. Influence of fast pyrolysis temperature on biochar labile fraction and short-
305 term carbon loss in a loamy soil. *Biomass and Bioenergy* 35, 1182-1189.
- 306 Bucheli, T.D., Blum, F., Desaulles, A., Gustafsson, Ö., 2004. Polycyclic aromatic hydrocarbons,
307 black carbon, and molecular markers in soils of Switzerland. *Chemosphere* 56, 1061-1076.
- 308 Cohen-Ofri, I., Weiner, L., Boaretto, E., Mintz, G., Weiner, S., 2006. Modern and fossil charcoal:
309 aspects of structure and diagenesis. *Journal Of Archaeological Science* 33, 428-439.
- 310 Colmsjö, A., 1998. Concentration and Extraction of PAHs from Environmental Samples, in:
311 Neilson, A. (Ed.), *PAHs and Related Compounds*. Springer, Berlin, Germany, pp. 55-76.
- 312 Coppola, A.I., Ziolkowski, L.A., Druffel, E.R.M., 2013. Extraneous carbon assessments in
313 radiocarbon measurements of black carbon in environmental matrices. *Radiocarbon* 55, 1631-
314 1640.
- 315 Denis, E.H., Toney, J.L., Tarozo, R., Scott Anderson, R., Roach, L.D., Huang, Y., 2012.
316 Polycyclic aromatic hydrocarbons (PAHs) in lake sediments record historic fire events: Validation
317 using HPLC-fluorescence detection. *Organic Geochemistry* 45, 7-17.
- 318 Dittmar, T., 2008. The molecular level determination of black carbon in marine dissolved organic
319 matter. *Organic Geochemistry* 39, 396-407.
- 320 Fabbri, D., Rombolà, A.G., Torri, C., Spokas, K.A., 2013. Determination of polycyclic aromatic
321 hydrocarbons in biochar and biochar amended soil. *Journal of Analytical and Applied Pyrolysis*
322 103, 60-67.

323 Fang, Y., Singh, B., Singh, B.P., Krull, E., 2014. Biochar carbon stability in four contrasting soils.
324 *European Journal of Soil Science* 65, 60-71.

325 Flannigan, M., Cantin, A.S., De Groot, W.J., Wotton, M., Newbery, A., Gowman, L.M., 2013.
326 Global wildland fire season severity in the 21st century. *Forest Ecology and Management* 294,
327 54-61.

328 Franklin, R.E., 1951. Crystallite growth in graphitizing and non-graphitizing carbons.
329 *Proceedings of the Royal Society of London Series A* 209, 196-218.

330 Gierga, M., Schneider, M.P.W., Wiedemeier, D.B., Lang, S.Q., Smittenberg, R.H., Hajdas, I.,
331 Bernasconi, S., Schmidt, M.W.I., 2014. Purification of fire-derived markers for μg scale isotope
332 analysis ($\delta^{13}\text{C}$, $\Delta^{14}\text{C}$) using high-performance liquid chromatography (HPLC). *Organic*
333 *Geochemistry* (in press).

334 Glaser, B., Haumaier, L., Guggenberger, G., Zech, W., 1998. Black carbon in soils: the use of
335 benzenecarboxylic acids as specific markers. *Organic Geochemistry* 29, 811-819.

336 Guggenberger, G., Rodionov, A., Shibistova, O., Grabe, M., Kasansky, O.A., Fuchs, H.,
337 Mikheyeva, N., Zrazhevskaya, G., Flessa, H., 2008. Storage and mobility of black carbon in
338 permafrost soils of the forest tundra ecotone in Northern Siberia. *Global Change Biology* 14,
339 1367-1381.

340 Hale, S.E., Lehmann, J., Rutherford, D., Zimmerman, A.R., Bachmann, R.T., Shitumbanuma, V.,
341 O'Toole, A., Sundqvist, K.L., Arp, H.P.H., Cornelissen, G., 2012. Quantifying the total and
342 bioavailable polycyclic aromatic hydrocarbons and dioxins in biochars. *Environmental Science &*
343 *Technology* 46, 2830-2838.

344 Hammes, K., Schmidt, M.W.I., Smernik, R.J., Currie, L.A., Ball, W.P., Nguyen, T.H.,
345 Louchouart, P., Houel, S., Gustafsson, O., Elmquist, M., Cornelissen, G., Skjemstad, J.O.,
346 Masiello, C.A., Song, J., Peng, P., Mitra, S., Dunn, J.C., Hatcher, P.G., Hockaday, W.C., Smith,
347 D.M., Hartkopf-Froeder, C., Boehmer, A., Luer, B., Huebert, B.J., Amelung, W., Brodowski, S.,
348 Huang, L., Zhang, W., Gschwend, P.M., Flores-Cervantes, D.X., Largeau, C., Rouzaud, J.N.,
349 Rumpel, C., Guggenberger, G., Kaiser, K., Rodionov, A., Gonzalez-Vila, F.J., Gonzalez-Perez,
350 J.A., de la Rosa, J.M., Manning, D.A.C., Lopez-Capel, E., Ding, L., 2007. Comparison of
351 quantification methods to measure fire-derived (black/elemental) carbon in soils and sediments
352 using reference materials from soil, water, sediment and the atmosphere. *Global*
353 *Biogeochemical Cycles* 21, GB3016.

354 Hammes, K., Torn, M., Lapenas, A., Schmidt, M., 2008. Centennial black carbon turnover
355 observed in a Russian steppe soil. *Biogeosciences* 5, 1339-1350.

356 Harvey, O., Kuo, L.-J., Zimmerman, A., Louchouart, P., Amonette, J., Herbert, B., 2012a. An
357 index-based approach to assessing recalcitrance and soil carbon sequestration potential of
358 engineered black carbons (biochars). *Environmental Science & Technology* 46, 1415-1421.

359 Harvey, O.R., Herbert, B.E., Kuo, L.-J., Louchouart, P., 2012b. Generalized two-dimensional
360 perturbation correlation infrared spectroscopy reveals mechanisms for the development of
361 surface charge and recalcitrance in plant-derived biochars. *Environmental Science &*
362 *Technology* 46, 10641-10650.

363 Hilber, I., Blum, F., Leifeld, J., Schmidt, H.P., Bucheli, T.D., 2012. Quantitative determination of
 364 PAHs in biochar: A prerequisite to ensure its quality and safe application. *Journal of Agricultural*
 365 *and Food Chemistry* 60, 3042-3050.

366 Keiluweit, M., Kleber, M., Sparrow, M.A., Simoneit, B.R.T., Prah, F.G., 2012. Solvent-
 367 extractable polycyclic aromatic hydrocarbons in biochar: influence of pyrolysis temperature and
 368 feedstock. *Environmental Science & Technology* 46, 9333-9341.

369 Keiluweit, M., Nico, P.S., Johnson, M.G., Kleber, M., 2010. Dynamic molecular structure of plant
 370 biomass-derived black carbon (biochar). *Environmental Science & Technology* 44, 1247-1253.

371 Kelly, R., Chipman, M.L., Higuera, P.E., Stefanova, I., Brubaker, L.B., Hu, F.S., 2013. Recent
 372 burning of boreal forests exceeds fire regime limits of the past 10,000 years. *Proceedings of the*
 373 *National Academy of Sciences of the United States of America* 110, 13055-13060.

374 Kloss, S., Zehetner, F., Dellantonio, A., Hamid, R., Ottner, F., Liedtke, V., Schwanninger, M.,
 375 Gerzabek, M.H., Soja, G., 2012. Characterization of slow pyrolysis biochars: Effects of
 376 feedstocks and pyrolysis temperature on biochar properties. *Journal of Environmental Quality*
 377 41, 990-1000.

378 Kuzyakov, Y., Bogomolova, I., Glaser, B., 2014. Biochar stability in soil: Decomposition during
 379 eight years and transformation as assessed by compound-specific ¹⁴C analysis. *Soil Biology*
 380 *and Biochemistry* 70, 229-236.

381 Lehmann, J., Joseph, S., 2009. Biochar for Environmental Management - An Introduction, in:
 382 Lehmann, J., Joseph, S. (Eds.), *Biochar for Environmental Management: Science and*
 383 *Technology*. Earthscan, London, UK, pp. 1-12.

384 Marschner, B., Brodowski, S., Dreves, A., Gleixner, G., Gude, A., Grootes, P.M., Hamer, U.,
 385 Heim, A., Jandl, G., Ji, R., Kaiser, K., Kalbitz, K., Kramer, C., Leinweber, P., Rethemeyer, J.,
 386 Schaeffer, A., Schmidt, M.W.I., Schwark, L., Wiesenberger, G.L.B., 2008. How relevant is
 387 recalcitrance for the stabilization of organic matter in soils? *Journal of Plant Nutrition and Soil*
 388 *Science* 171, 91-110.

389 Masiello, C.A., 2004. New directions in black carbon organic geochemistry. *Marine Chemistry*
 390 92, 201-213.

391 McBeath, A., Smernik, R., Plant, E., 2011. Determination of the aromaticity and the degree of
 392 aromatic condensation of a thermosequence of wood charcoal using NMR. *Organic*
 393 *Geochemistry* 42, 1194-1202.

394 McGrath, T.E., Geoffrey Chan, W., Hajaligol, M.R., 2003. Low temperature mechanism for the
 395 formation of polycyclic aromatic hydrocarbons from the pyrolysis of cellulose. *Journal of*
 396 *Analytical and Applied Pyrolysis* 66, 51-70.

397 Meyer, S., Glaser, B., Quicker, P., 2011. Technical, economical, and climate-related aspects of
 398 biochar production technologies: a literature review. *Environmental Science & Technology* 45,
 399 9473-9483.

400 Oleszczuk, P., Joško, I., Kuśmierz, M., 2013. Biochar properties regarding to contaminants
 401 content and ecotoxicological assessment. *Journal of Hazardous materials* 260, 375-382.

402 Oros, D.R., Simoneit, B.R.T., 2001a. Identification and emission factors of molecular tracers in
 403 organic aerosols from biomass burning Part 1. Temperate climate conifers. *Applied*
 404 *Geochemistry* 16, 1513-1544.

405 Oros, D.R., Simoneit, B.R.T., 2001b. Identification and emission factors of molecular tracers in
 406 organic aerosols from biomass burning Part 2. Deciduous trees. *Applied Geochemistry* 16,
 407 1545-1565.

408 Quilliam, R.S., Rangecroft, S., Emmett, B.A., Deluca, T.H., Jones, D.L., 2013. Is biochar a
 409 source or sink for polycyclic aromatic hydrocarbon (PAH) compounds in agricultural soils? *GCB*
 410 *Bioenergy* 5, 96-103.

411 R, 2011. R: A language and environment for statistical computing. R Foundation for Statistical
 412 Computing.

413 Radke, M., Willsch, H., Welte, D.H., 1980. Preparative hydrocarbon group type determination by
 414 automated medium pressure liquid chromatography. *Analytical Chemistry* 52, 406-411.

415 Rennert, T., Kaufhold, S., Brodowski, S., Mansfeldt, T., 2008. Interactions of ferricyanide with
 416 humic soils and charred straw. *European Journal of Soil Science* 59, 348-358.

417 Rodionov, A., Amelung, W., Peinemann, N., Haumaier, L., Zhang, X., Kleber, M., Glaser, B.,
 418 Urusevskaya, I., Zech, W., 2010. Black carbon in grassland ecosystems of the world. *Global*
 419 *Biogeochemical Cycles* 24, GB3013.

420 Sánchez-García, L., de Andrés, J.R., Gélinas, Y., Schmidt, M.W.I., Louchouart, P., 2013.
 421 Different pools of black carbon in sediments from the Gulf of Cádiz (SW Spain): Method
 422 comparison and spatial distribution. *Marine Chemistry* 151, 13-22.

423 Schmidt, M.W.I., Noack, A.G., 2000. Black carbon in soils and sediments: Analysis, distribution,
 424 implications, and current challenges. *Global Biogeochemical Cycles* 14, 777-793.

425 Schmidt, M.W.I., Skjemstad, J.O., Czimczik, C.I., Glaser, B., Prentice, K.M., Gelinas, Y.,
 426 Kuhlbusch, T.A.J., 2001. Comparative analysis of black carbon in soils. *Global Biogeochemical*
 427 *Cycles* 15, 163-167.

428 Schneider, M.P.W., Hilf, M., Vogt, U.F., Schmidt, M.W.I., 2010. The benzene polycarboxylic acid
 429 (BPCA) pattern of wood pyrolyzed between 200 °C and 1000 °C. *Organic Geochemistry* 41,
 430 1082-1088.

431 Schneider, M.P.W., Smittenberg, R., Dittmar, T., Schmidt, M.W.I., 2011. Comparison of gas with
 432 liquid chromatography for the determination of benzenepolycarboxylic acids as molecular tracers
 433 of black carbon. *Organic Geochemistry* 42, 275-282.

434 Simoneit, B.R.T., 2002. Biomass burning -- a review of organic tracers for smoke from
 435 incomplete combustion. *Applied Geochemistry* 17, 129-162.

436 Simoneit, B.R.T., Fraser, M.P., Schauer, J.J., Nolte, C.G., Oros, D.R., Elias, V.O., Rogge, W.F.,
 437 Cass, G.R., 1999. Levoglucosan, a tracer for cellulose in biomass burning and atmospheric
 438 particles. *Atmospheric Environment* 33, 173-182.

439 Slater, G.F., Benson, A.A., Marvin, C., Muir, D., 2013. PAH fluxes to Siskiwit revisited: Trends in
 440 fluxes and sources of pyrogenic PAH and perylene constrained via radiocarbon analysis.
 441 *Environmental Science and Technology* 47, 5066-5073.

442 Wiedemeier, D.B., Abiven, S., Hockaday, W.C., Keiluweit, M., Kleber, M., Pyle, L.A., Masiello,
 443 C.A., McBeath, A.V., Nico, P.S., Schneider, M.P.W., Smernik, R.J., Wiesenberg, G.L.B.,
 444 Schmidt, M.W.I., 2014. Aromaticity and aromatic condensation of chars. *Organic Geochemistry*
 445 (in review).

446 Wiedemeier, D.B., Hilf, M.D., Smittenberg, R.H., Haberle, S.G., Schmidt, M.W.I., 2013. Improved
 447 assessment of pyrogenic carbon quantity and quality in environmental samples by high-
 448 performance liquid chromatography. *Journal of Chromatography A* 1304, 246-250.

449 Wiesenberg, G.L.B., Brodowski, S., 2007. Veränderungen der Zusammensetzung und
 450 Überlagerung von Lipiden und Black Carbon infolge der Verkohlung von Pflanzenmaterial.
 451 *Mitteilungen der Deutschen Bodenkundlichen Gesellschaft* 110, 315-316.

452 Wiesenberg, G.L.B., Lehndorff, E., Schwark, L., 2009. Thermal degradation of rye and maize
 453 straw: lipid pattern changes as a function of temperature. *Organic Geochemistry* 40, 167-174.

454 Wiesenberg, G.L.B., Schmidt, M.W.I., Schwark, L., 2008. Plant and soil lipid modifications under
 455 elevated atmospheric CO₂ conditions: I. Lipid distribution patterns. *Organic Geochemistry* 39,
 456 91-102.

457 Wiesenberg, G.L.B., Schwark, L., Schmidt, M.W.I., 2004. Improved automated extraction and
 458 separation procedure for soil lipid analyses. *European Journal of Soil Science* 55, 349-356.

459 Wilcke, W., 2000. Polycyclic aromatic hydrocarbons (PAHs) in soil — A review. *Journal of Plant*
 460 *Nutrition and Soil Science* 163, 229-248.

461 Wurster, C.M., Saiz, G., Schneider, M.P.W., Schmidt, M.W.I., Bird, M.I., 2013. Quantifying
 462 pyrogenic carbon from thermosequences of wood and grass using hydrogen pyrolysis. *Organic*
 463 *Geochemistry* 62, 28-32.

464 Xu, L., Zheng, M., Ding, X., Edgerton, E.S., Reddy, C.M., 2012. Modern and fossil contributions
 465 to polycyclic aromatic hydrocarbons in PM 2.5 from North Birmingham, Alabama in the
 466 Southeastern U.S. *Environmental Science and Technology* 46, 1422-1429.

467 Yunker, M.B., Macdonald, R.W., Vingarzan, R., Mitchell, R.H., Goyette, D., Sylvestre, S., 2002.
 468 PAHs in the Fraser River basin: a critical appraisal of PAH ratios as indicators of PAH source
 469 and composition. *Organic Geochemistry* 33, 489-515.

470 Ziolkowski, L., Druffel, E., 2010. Aged black carbon identified in marine dissolved organic
 471 carbon. *Geophysical Research Letters* 37, L16601.

472

473

474